

Is π Normal?

David H. Bailey

Lawrence Berkeley National Laboratory

Berkeley, CA, USA 94720

`dhbailey@lbl.gov`

Collaborators:

Jonathan M. Borwein, Simon Fraser University, Canada

Peter B. Borwein, Simon Fraser University, Canada

David J. Broadhurst, Open University, UK

Richard E. Crandall, Center for Advanced Computation, Reed College

Helaman R. P. Ferguson, Center for Computing Sciences

The First 1000 Decimal Digits of Pi

3.

1415926535897932384626433832795028841971693993751058209749445923078164062862089986280348253421170679
8214808651328230664709384460955058223172535940812848111745028410270193852110555964462294895493038196
4428810975665933446128475648233786783165271201909145648566923460348610454326648213393607260249141273
7245870066063155881748815209209628292540917153643678925903600113305305488204665213841469519415116094
3305727036575959195309218611738193261179310511854807446237996274956735188575272489122793818301194912
9833673362440656643086021394946395224737190702179860943702770539217176293176752384674818467669405132
0005681271452635608277857713427577896091736371787214684409012249534301465495853710507922796892589235
4201995611212902196086403441815981362977477130996051870721134999999837297804995105973173281609631859
5024459455346908302642522308253344685035261931188171010003137838752886587533208381420617177669147303
5982534904287554687311595628638823537875937519577818577805321712268066130019278766111959092164201989

The First 1000 Hexadecimal (Base 16) Digits of Pi

3.

243f6a8885a308d313198a2e03707344a093822299f31d0082efa98ec4e6c89452821e638d01377be5466cf34e90c6cc0
ac29b7c97c50dd3f84d5b5b54709179216d5d98979fb1bd1310ba698dfb5ac2ffd72dbd01adfb7b8e1afed6a267e96ba7c
9045f12c7f9924a19947b3916cf70801f2e2858efc16636920d871574e69a458fea3f4933d7e0d95748f728eb658718bcd
5882154aee7b54a41dc25a59b59c30d5392af26013c5d1b023286085f0ca417918b8db38ef8e79dc b0603a180e6c9e0e8b
b01e8a3ed71577c1bd314b2778af2fda55605c60e65525f3aa55ab945748986263e8144055ca396a2aab10b6b4cc5c3411
41e8cea15486af7c72e993b3ee1411636fbc2a2ba9c55d741831f6ce5c3e169b87931eafdb6ba336c24cf5c7a3253812895
86773b8f48986b4bb9afc4bfe81b6628219361d809ccfb21a991487cac605dec8032ef845d5de98575b1dc262302eb651b
8823893e81d396acc50f6d6ff383f442392e0b4482a484200469c8f04a9e1f9b5e21c66842f6e96c9a670c9c61abd388f0
6a51a0d2d8542f68960fa728ab5133a36eef0b6c137a3be4ba3bf0507efb2a98a1f1651d39af017666ca593e82430e888c
ee8619456f9fb47d84a5c33b8b5eb06f75d885c12073401a449f56c16aa64ed3aa62363f77061bfedf72429b023d37d0

On-line tool searches for any pattern in the first four billion digits of π :
<http://pi.nersc.gov>

Normality

The real number α is *normal* to base b if every sequence of m digits in the base- b expansion of α appears with limiting frequency b^{-m} .

Almost all real numbers are normal (from measure theory). Widely believed to be normal base b for all bases b :

- π and e .
- $\log 2$ and $\sqrt{2}$.
- The golden mean $\tau = (1 + \sqrt{5})/2$.
- *Every* irrational algebraic number.
- Many other “natural” irrational constants.

But there are *no* proofs for any of these constants, for any base. Normality proofs exist only for handful of artificially constructed constants, such as Champernowne’s number: 0.1234567891011121314...

Integer Relation Detection

Given a real or complex vector $x = (x_1, x_2, \dots, x_n)$ an *integer relation* (IR) algorithm seeks integers a_i , not all zero, such that

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = 0$$

to within the available numerical accuracy.

- Original IR algorithm found in 1977 by Helaman Ferguson and Rodney Forcade.
- Current state of art: Ferguson’s “PSLQ” algorithm — recently named one of ten “algorithms of the century” by *Computing in Science and Engineering*.
- Very high numeric precision (hundreds or thousands of digits) must be employed in integer relation calculations.

Applications of PSLQ: Recognizing Numeric Constants

If α is algebraic of degree n , the polynomial satisfied by α can be found by computing the vector $(1, \alpha, \alpha^2, \dots, \alpha^n)$ to high precision, and then applying PSLQ.

Example:

Let $B_3 = 3.54409035955 \dots$ be the third bifurcation point of the logistic map $x_{k+1} = rx_k(1 - x_k)$. In other words, B_3 is the smallest r such that successive iterates x_k exhibit eight-way periodicity instead of four-way periodicity.

Computations using a predecessor algorithm to PSLQ found that B_3 is a root of the polynomial

$$0 = 4913 + 2108t^2 - 604t^3 - 977t^4 + 8t^5 + 44t^6 + 392t^7 - 193t^8 - 40t^9 + 48t^{10} - 12t^{11} + t^{12}$$

Recently a PSLQ program found that $\alpha = -B_4(B_4 - 2)$ satisfies a 120-degree polynomial, so that B_4 satisfies a 240-degree polynomial.

Applications of PSLQ: Euler Sums

Let $\zeta(t) = \sum_{j=1}^{\infty} j^{-t}$ be the Riemann zeta function, and $\text{Li}_n(x) = \sum_{j=1}^{\infty} x^j j^{-n}$ the polylogarithm function. The following were found using PSLQ computations:

$$\begin{aligned}
\sum_{k=1}^{\infty} \left(1 + \frac{1}{2} + \cdots + \frac{1}{k}\right)^2 (k+1)^{-4} &= \frac{37}{22680} \pi^6 - \zeta^2(3) \\
\sum_{k=1}^{\infty} \left(1 + \frac{1}{2} + \cdots + \frac{1}{k}\right)^3 (k+1)^{-6} &= \zeta^3(3) + \frac{197}{24} \zeta(9) + \frac{1}{2} \pi^2 \zeta(7) \\
&\quad - \frac{11}{120} \pi^4 \zeta(5) - \frac{37}{7560} \pi^6 \zeta(3) \\
\sum_{k=1}^{\infty} \left(1 - \frac{1}{2} + \cdots + (-1)^{k+1} \frac{1}{k}\right)^2 (k+1)^{-3} &= 4 \text{Li}_5\left(\frac{1}{2}\right) - \frac{1}{30} \ln^5(2) - \frac{17}{32} \zeta(5) \\
&\quad - \frac{11}{720} \pi^4 \ln(2) + \frac{7}{4} \zeta(3) \ln^2(2) \\
&\quad + \frac{1}{18} \pi^2 \ln^3(2) - \frac{1}{8} \pi^2 \zeta(3)
\end{aligned}$$

Applications of PSLQ: Apery Sums

It has been known for some time, through the research of Apery, that

$$\begin{aligned}\zeta(2) &= 3 \sum_{k=1}^{\infty} \frac{1}{k^2 \binom{2k}{k}} \\ \zeta(3) &= \frac{5}{2} \sum_{k=1}^{\infty} \frac{(-1)^{k-1}}{k^3 \binom{2k}{k}} \\ \zeta(4) &= \frac{36}{17} \sum_{k=1}^{\infty} \frac{1}{k^4 \binom{2k}{k}}\end{aligned}$$


These results have led many to suggest that

$$S(n) = \sum_{k>0} \frac{1}{k^n \binom{2k}{k}},$$

for $n > 4$, might be a simple constant. It has now been shown that $S(n)$ can be expressed in terms of the Riemann zeta function $\zeta(n)$ and Clausen's function $M(a, b)$. A sample evaluation is

$$\begin{aligned}S(9) &= \pi \left[2M(7, 1) + \frac{8}{3}M(5, 3) + \frac{8}{9}\zeta(2)M(5, 1) \right] - \frac{13921}{216}\zeta(9) \\ &\quad + \frac{6211}{486}\zeta(7)\zeta(2) + \frac{8101}{648}\zeta(6)\zeta(3) + \frac{331}{18}\zeta(5)\zeta(4) - \frac{8}{9}\zeta^3(3)\end{aligned}$$

Ten Tetrahedral Cases from Quantum Field Theory



Evaluations of constants associated with the ten cases:

$$V_1 = 6\zeta(3) + 3\zeta(4)$$

$$U = \sum_{j>k>0} \frac{(-1)^{j+k}}{j^3 k}$$

$$V_{2A} = 6\zeta(3) - 5\zeta(4)$$

$$C = \sum_{k>0} \sin(\pi k/3)/k^2$$

$$V_{2N} = 6\zeta(3) - \frac{13}{2}\zeta(4) - 8U$$

$$V = \sum_{j>k>0} (-1)^j \cos(2\pi k/3)/(j^3 k)$$

$$V_{3T} = 6\zeta(3) - 9\zeta(4)$$

$$V_{3S} = 6\zeta(3) - \frac{11}{2}\zeta(4) - 4C^2$$

$$V_{3L} = 6\zeta(3) - \frac{15}{4}\zeta(4) - 6C^2$$

$$V_{4A} = 6\zeta(3) - \frac{77}{12}\zeta(4) - 6C^2$$

$$V_{4N} = 6\zeta(3) - 14\zeta(4) - 16U$$

$$V_5 = 6\zeta(3) - \frac{469}{27}\zeta(4) + \frac{8}{3}C^2 - 16V$$

$$V_6 = 6\zeta(3) - 13\zeta(4) - 8U - 4C^2$$

Peter Borwein's Observation on the Binary Digits of $\log 2$

In 1995, Peter Borwein observed that an individual binary digit of $\log 2$ can be calculated by using a very simple algorithm:

Let $\{\cdot\}$ denote the fractional part. Then we can write

$$\begin{aligned} \{2^d \log 2\} &= \left\{ 2^d \sum_{k=1}^{\infty} \frac{1}{k \cdot 2^k} \right\} = \left\{ \sum_{k=1}^{\infty} \frac{2^{d-k}}{k} \right\} \\ &= \left\{ \sum_{k=1}^d \frac{2^{d-k}}{k} \right\} + \left\{ \sum_{k=d+1}^{\infty} \frac{2^{d-k}}{k} \right\} \\ &= \left\{ \sum_{k=1}^d \frac{2^{d-k} \bmod k}{k} \right\} + \left\{ \sum_{k=d+1}^{\infty} \frac{2^{d-k}}{k} \right\} \end{aligned}$$

- The numerators $2^{d-k} \bmod k$ can be very rapidly evaluated using the binary algorithm for exponentiation performed modulo k .
- Only a few terms of the second summation need be evaluated.
- All computations can be done with ordinary 64-bit floating-point arithmetic.

A More General Result

Any constant α given by a formula of the type

$$\alpha = \sum_{k=0}^{\infty} \frac{p(k)}{b^k q(k)}$$

(where $p(k)$ and $q(k)$ are integer polynomials, $\deg p < \deg q$ and q has no zeroes for positive k) has the rapid individual digit computation property.

Is there a formula of this type for π ? None was known in 1995.

The BBP Formula for π

By applying DHB's PSLQ computer program to set of computed constants for which formulas of this type were known, with the numerical value of π appended, Simon Plouffe found this formula for π :

$$\pi = \sum_{k=0}^{\infty} \frac{1}{16^k} \left(\frac{4}{8k+1} - \frac{2}{8k+4} - \frac{1}{8k+5} - \frac{1}{8k+6} \right)$$

Question: Why wasn't this formula discovered 250 years ago?

Proof of the BBP Pi Formula

We can write

$$\int_0^{1/\sqrt{2}} \frac{x^{j-1} dx}{1-x^8} = \int_0^{1/\sqrt{2}} \sum_{k=0}^{\infty} x^{8k+j-1} dx = \frac{1}{2^{j/2}} \sum_{k=0}^{\infty} \frac{1}{16^k(8k+j)}$$

Thus

$$\begin{aligned} & \sum_{k=0}^{\infty} \frac{1}{16^k} \left(\frac{4}{8k+1} - \frac{2}{8k+4} - \frac{1}{8k+5} - \frac{1}{8k+6} \right) \\ &= \int_0^{1/\sqrt{2}} \left(\frac{4\sqrt{2} - 8x^3 - 4\sqrt{2}x^4 - 8x^5}{1-x^8} \right) dx \\ &= \int_0^1 \frac{16(4-2y^3-y^4-y^5) dy}{16-y^8} \\ &= \int_0^1 \frac{16(y-1) dy}{(y^2-2)(y^2-2y+2)} \\ &= \int_0^1 \frac{4y dy}{y^2-2} - \int_0^1 \frac{(4y-8) dy}{y^2-2y+2} \\ &= \pi \end{aligned}$$

The BBP Algorithm for Computing Individual Hex Digits of Pi

Let S_1 be the first of the four sums in the formula for π .

$$\begin{aligned} (16^n S_1) \bmod 1 &= \left(\sum_{k=0}^{\infty} \frac{16^{n-k}}{8k+1} \right) \bmod 1 = \left(\sum_{k=0}^n \frac{16^{n-k}}{8k+1} + \sum_{k=n+1}^{\infty} \frac{16^{n-k}}{8k+1} \right) \bmod 1 \\ &= \left(\sum_{k=0}^n \frac{16^{n-k} \bmod 8k+1}{8k+1} + \sum_{k=n+1}^{\infty} \frac{16^{n-k}}{8k+1} \right) \bmod 1 \end{aligned}$$

1. Compute each numerator of each term in the first sum using the binary algorithm for exponentiation, reducing each product modulo $8k+1$.
2. Divide each numerator by its respective denominator $8k+1$.
3. Sum the terms of the first series, discarding integer parts.
4. Compute the second sum (just a few terms are needed).
5. Add the two sum results, again discarding the integer part.
6. Repeat for S_1, S_2, S_3, S_4 , and calculate $4S_1 - 2S_2 - S_3 - S_4$.
7. The resulting fraction, when expressed in hexadecimal format, gives the first few hex digits of π beginning at position $n+1$.

Ordinary 64-bit or 128-bit floating-point arithmetic suffices for these operations — multiple precision arithmetic software is *not* required.

Some Computational Results

Position	Hex Digits of π Starting at Position
10^6	26C65E52CB4593
10^7	17AF5863EFFED8D
10^8	ECB840E21926EC
10^9	85895585A0428B
10^{10}	921C73C6838FB2
10^{11}	9C381872D27596
1.25×10^{12}	[1] 07E45733CC790B
2.5×10^{14}	[2] E6216B069CB6C1

[1] Babrice Bellard, France, 1999
 [2] Colin Percival, Canada, 2000

Are There BBP-Type Formulas for Pi in Other Bases?

Jonathan Borwein, David Borwein and William Galway have now shown that there are no formulas of the type

$$\pi = \sum_{k=0}^{\infty} \frac{p(k)}{b^k q(k)}$$

(where $p(k)$ and $q(k)$ are integer polynomials, $\deg p < \deg q$ and q has no zeroes for positive k), except for $b = 2^r$ for some integer r .

Thus 16 can be thought of as the “natural” base for π .

Some Other Constants with Base 2 BBP-Type Formulas

$$\begin{aligned}
\log 3 &= \sum_{k=0}^{\infty} \frac{1}{4^k(2k+1)} \\
\log 7 &= \frac{3}{4} \sum_{k=0}^{\infty} \frac{1}{8^k} \left(\frac{2}{8k+1} + \frac{1}{8k+2} \right) \\
\pi^2 &= \frac{1}{8} \sum_{k=0}^{\infty} \frac{1}{64^k} \left(\frac{144}{(6k+1)^2} - \frac{216}{(6k+2)^2} - \frac{72}{(6k+3)^2} - \frac{54}{(6k+4)^2} + \frac{9}{(6k+5)^2} \right) \\
\log^2 2 &= \frac{1}{6} \sum_{k=0}^{\infty} \frac{1}{16^k} \left(\frac{16}{(8k+1)^2} - \frac{40}{(8k+2)^2} - \frac{8}{(8k+3)^2} - \frac{28}{(8k+4)^2} \right. \\
&\quad \left. - \frac{4}{(8k+5)^2} - \frac{10}{(8k+6)^2} + \frac{2}{(8k+7)^2} - \frac{3}{(8k+8)^2} \right) \\
\pi^2 - 6 \log^2 2 &= \frac{12}{\sum_{k=1}^{\infty} \frac{1}{k^2 2^k}} \\
\pi \sqrt{3} &= \frac{9}{32} \sum_{k=0}^{\infty} \frac{1}{64^k} \left(\frac{16}{6k+1} - \frac{8}{6k+2} - \frac{2}{6k+4} - \frac{1}{6k+5} \right)
\end{aligned}$$

An Arctan Formula

$$\tan^{-1}\left(\frac{4}{5}\right) = \frac{1}{2^{17}} \sum_{k=0}^{\infty} \frac{1}{2^{20k}} \left(\frac{524288}{40k+2} - \frac{393216}{40k+4} - \frac{491520}{40k+5} + \frac{163840}{40k+8} \right. \\ \left. + \frac{32768}{40k+10} - \frac{24576}{40k+12} + \frac{5120}{40k+15} + \frac{10240}{40k+16} + \frac{2048}{40k+18} \right. \\ \left. + \frac{1024}{40k+20} + \frac{640}{40k+24} + \frac{480}{40k+25} + \frac{128}{40k+26} - \frac{96}{40k+28} \right. \\ \left. + \frac{40}{40k+32} + \frac{8}{40k+34} - \frac{5}{40k+35} - \frac{6}{40k+36} \right)$$

Similar formulas have been found for arctans of numerous other rational arguments.

Some Base 3 BBP-Type Formulas

$$\begin{aligned}
\log 2 &= \frac{2}{27} \sum_{k=0}^{\infty} \frac{1}{81^k} \left(\frac{9}{4k+1} + \frac{1}{4k+3} \right) \\
&= \sum_{n=0}^{\infty} \frac{1}{9^n(2n-1)} \\
\pi^2 &= \frac{2}{27} \sum_{k=0}^{\infty} \frac{1}{729^k} \left(\frac{243}{(12k+1)^2} - \frac{405}{(12k+2)^2} - \frac{81}{(12k+4)^2} - \frac{27}{(12k+5)^2} \right. \\
&\quad \left. - \frac{72}{(12k+6)^2} - \frac{9}{(12k+7)^2} - \frac{9}{(12k+8)^2} - \frac{5}{(12k+10)^2} + \frac{1}{(12k+11)^2} \right) \\
6\sqrt{3} \tan^{-1} \left(\frac{\sqrt{3}}{7} \right) &= \sum_{k=0}^{\infty} \frac{1}{27^k} \left(\frac{3}{3k+1} + \frac{1}{3k+2} \right)
\end{aligned}$$

A Base 5 BBP-Type Formula

$$\frac{25}{2} \log \left(\frac{781}{256} \left(\frac{57 - 5\sqrt{5}}{57 + 5\sqrt{5}} \right)^{\sqrt{5}} \right) = \sum_{k=0}^{\infty} \frac{1}{5^{5k}} \left(\frac{5}{5k+2} + \frac{1}{5k+3} \right)$$

Two Base 10 BBP-Type Formulas

$$\begin{aligned} \log \left(\frac{9}{10} \right) &= - \sum_{k=1}^{\infty} \frac{1}{k10^k} \\ \log \left(\frac{1111111111}{387420489} \right) &= 10^{-8} \sum_{k=0}^{\infty} \frac{1}{10^{10k}} \left(\frac{10^8}{10k+1} + \frac{10^7}{10k+2} + \dots + \frac{1}{10k+9} \right) \end{aligned}$$

A Connection Between BBP-Type Formulas and Normality

Theorem: The BBP-type constant

$$\alpha = \sum_{k=0}^{\infty} \frac{p(k)}{b^k q(k)}$$

(where $p(k)$ and $q(k)$ are integer polynomials, $\deg p < \deg q$ and q has no zeroes for positive k) is normal base b if and only if the sequence $x_0 = 0$, and

$$x_n = \left(bx_{n-1} + \frac{p(n)}{q(n)} \right) \bmod 1$$

is equidistributed in the unit interval.

Proof Sketch: Let α_n be the base- b expansion of α after the n -th digit. Following the BBP approach, we can write

$$\begin{aligned} \alpha_n &= \left\{ \sum_{k=0}^n \frac{b^{n-k} p(k)}{q(k)} \right\} + \left\{ \sum_{k=n+1}^{\infty} \frac{b^{n-k} p(k)}{q(k)} \right\} \\ &= \left(b\alpha_{n-1} + \frac{p(n)}{q(n)} \right) \bmod 1 + E_n \end{aligned}$$

where E_n goes to zero.

Two Examples

1. Let $x_0 = 0$, and

$$x_n = \left(2x_{n-1} + \frac{1}{n} \right) \bmod 1$$

Is (x_n) equidistributed in $[0, 1)$?

2. Let $x_0 = 0$ and

$$x_n = \left(16x_{n-1} + \frac{120n^2 - 89n + 16}{512n^4 - 1024n^3 + 712n^2 - 206n + 21} \right) \bmod 1$$

Is (x_n) equidistributed in $[0, 1)$?

If answer to Question 1 is “yes”, then $\log 2$ is normal to base 2.

If answer to Question 2 is “yes”, then π is normal to base 16 (and hence to base 2 also).

Hypothesis A

Denote by $r_n = p(n)/q(n)$ a rational-polynomial function, $0 \leq \deg(p) < \deg(q)$. Let b be an integer, $b \geq 2$ and set $x_0 = 0$. Then the sequence

$$x_n = (bx_{n-1} + r_n) \bmod 1$$

either has a finite attractor or is equidistributed in $[0, 1)$.

Theorem: Assuming Hypothesis A, then any constant α given by a formula of the form

$$\alpha = \sum_{k=0}^{\infty} \frac{p(k)}{b^k q(k)}$$

(where $p(k)$ and $q(k)$ are integer polynomials, $\deg p < \deg q$ and q has no zeroes for positive k) is either normal base b or rational.

A Surprising Empirical Result

Recall the iteration associated with π : Let $x_0 = 0$ and

$$x_n = \left(16x_{n-1} + \frac{120n^2 - 89n + 16}{512n^4 - 1024n^3 + 712n^2 - 206n + 21} \right) \bmod 1$$

Let y_n be the integer sequence defined as the index of the 16 subintervals of the unit interval where x_n lies, i.e. $y_n = \lfloor 16x_n \rfloor$. Then

Conjecture: The sequence (y_n) is precisely the hexadecimal expansion of π .

This has been verified by computer to 100,000 places.

A Class of Provably Normal Constants

Using the BBP approach, Richard Crandall and DHB have now proven normality for a class of constants, the simplest instance of which is

$$\begin{aligned}\alpha_{2,3} &= \sum_{k=1}^{\infty} \frac{1}{3^k 2^{3^k}} \\ &= 0.041883680831502985071252898624571682426096 \dots_{10} \\ &= 0.0AB8FE38F684BDA12F684BF35BA781948B0FCD6E9E0 \dots_{16}.\end{aligned}$$

$\alpha_{2,3}$ was actually proven normal base 2 in a little-known paper by Stoneham in 1977. Crandall and DHB proved normality and transcendence for an uncountably infinite class that includes $\alpha_{2,3}$:

$$\alpha_{2,3}(r) = \sum_{k=1}^{\infty} \frac{1}{3^k 2^{3^k + r_k}}$$

where r_k is the k -th bit in the binary expansion of $r \in (0, 1)$.

These constants also possess the rapid individual digit computation property. The googol-th binary digit of $\alpha_{2,3}$ is zero.

The Sequence Associated with $\alpha_{2,3}$.

Let $x_0 = 0$, and define

$$\begin{aligned} x_n &= (2x_{n-1}) \bmod 1 && \text{for } n \neq 3^k \\ &= (2x_{n-1} + 1/n) \bmod 1 && \text{for } n = 3^k \end{aligned}$$

The sequence (x_n) is merely the concatenation of primitive linear congruential pseudorandom sequences, each of length $2 \cdot 3^k$:

0, repeated 3 times,

$\frac{1}{3}, \frac{2}{3}$, repeated 3 times,

$\frac{4}{9}, \frac{8}{9}, \frac{7}{9}, \frac{5}{9}, \frac{1}{9}, \frac{2}{9}$, repeated 3 times,

$\frac{13}{27}, \frac{26}{27}, \frac{25}{27}, \frac{23}{27}, \frac{19}{27}, \frac{11}{27}, \frac{22}{27}, \frac{17}{27}, \frac{7}{27}, \frac{14}{27}, \frac{1}{27}, \frac{2}{27}, \frac{4}{27}, \frac{8}{27}, \frac{16}{27}, \frac{5}{27}, \frac{10}{27}, \frac{20}{27},$

repeated 3 times, etc.

New Results for Irrational Algebraic Numbers

Theorem A: Let $B_n(\alpha)$ be the number of ones in the binary expansion of α . Then for any irrational algebraic number α ,

$$\liminf_{n \rightarrow \infty} \frac{B_n(\alpha)/n}{\log_2(n)/n} \geq 1$$

Theorem B: If α is the square root of an integer or rational number, then for some constant C ,

$$\liminf_{n \rightarrow \infty} \frac{B_n(\alpha)/n}{C/\sqrt{n}} \geq 1$$

This result can be extended to the largest real root of an m -th degree integer-coefficient polynomial, where \sqrt{n} is replaced with $n^{1/m}$.

For Full Details

- David H. Bailey, Peter B. Borwein and Simon Plouffe, “On The Rapid Computation of Various Polylogarithmic Constants,” *Mathematics of Computation*, vol. 66, no. 218, 1997, pp. 903–913.
- David H. Bailey, “A Compendium of BBP-Type Formulas,” 2002.
- David H. Bailey and Richard E. Crandall, “On the Random Character of Fundamental Constant Expansions,” *Experimental Mathematics*, June 2001.
- David H. Bailey and Richard E. Crandall, “Random Generators and Normal Numbers,” 2002.

These are available at:

<http://www.nersc.gov/~dhbailey/dhbpapers>